

LYGOSOMA BOWRINGII (Bowring's Supple Skink). BANGLADESH: COMILLA DISTRICT: Lalmai-Tipperah Hills (23°22'N, 91°07'E), 12 July 2006. S. Mahony and A.H.M. Ali Reza. Wildlife Laboratory, Department of Zoology, Jahangirnagar University, Savar, Dhaka (JU 0050). Photograph deposited at USDZ, Raffles Museum of Biodiversity Research, National University of Singapore (ZRC[IMG].2.1). Verified by Indraneil Das. First record for Bangladesh and western range extension from nearest populations in Myanmar (of which the border is > 240 km w of Comilla) (Das 2002. *A Photographic Guide to Snakes and Other Reptiles of India*. New Holland Publishers [UK] Ltd., London. 144 pp.). Supported by Cleveland Metroparks Zoo and Rufford Foundation to the second author, which partially supported the first author. Bangladesh Forest Department issued permission (CCF [Wildlife]/2M-47/2006) for this work to the second author. Thanks are due to Md. Kamal Hossain from JU for his enthusiastic assistance in the field.

Submitted by **STEPHEN MAHONY**, 3 Park View Lawns, Greenpark, Clondalkin, Dublin 22, Ireland (e-mail: stephenmahony2@gmail.com); and **A. H. M. ALI REZA**, Department of Natural Resources Management, Texas Tech University, Lubbock, Texas 79409, USA (e-mail: ali.reza@ttu.edu).
